Neuroscience Letters, 124 (1991) 277–279
Elsevier Scientific Publishers Ireland Ltd.

AUTHOR INDEX

Abdullaev, Y.G., see Bechtereva, N.P., 61
Alessandri, M., see Tramontana, M., 79
Amador, M. and Dani, J.A., Protein kinase inhibitor, H-7, directly affects *N*-methyl-D-aspartate receptor channels, 251
Anderson, D.J., see Stemple, D.L., 57
Anderson, K.J., see Mitchell, J.J., 269
Arai, Y., see Matsumoto, A., 225
Araki, H., see Ishihara, A., 141
Aunis, D., see Jung, D., 87
Avoli, M., see Hwa, G.G.C., 65

Baba, A., see Koyama, Y., 235
Baimbridge, K.G., see Church, J., 232
Barrett, P., see Fraser, S.P., 242
Beart, P.M., Mercer, L.D. and Jarrot, B., [^{125}I]Ifenprodil: a convenient radioligandfor binding and autoradiographic studies of the polyamine-sensitive siteof the NMDA receptor, 187
Bechtereva, N.P., Abdullaev, Y.G. and Medvedev, S.V., Neuronal activity in frontal speech area 44 of the human cerebral cortex during word recognition, 61
Belan, P.V. and Osipenko, O., Blocking effect of La^{3+} ions on transmembrane ionic current evoked by intracellular cyclic AMP injection in identified *Helix pomatia* neurons, 137
Belhage, B., see Hansen, G.H., 162
Bennett, G.J., see Wakisaka, S., 200
Benoliel, J.J., Bourgoin, S., Mauborgne, A., Legrand, J.C., Hamon, M. and Cesselin, F., Differential inhibitory/stimulatory modulationof spinal CCK release by μ and δ opioid agonists, and selective blockade of μ-dependent inhibition by κ receptor stimulation, 204
Berod, A., see Kitabgi, P., 9
Berthold, C.-H., see Gatzinsky, K.P., 264
Beyreuther, K., see Prior, R., 69
Bianco, E.D., see Tramontana, M., 79
Birrell, G.J., McQueen, D.S., Iggo, A., Coleman, R.A. and Grubb, B.D., PGI_2-induced activation and sensitization of articular mechanonociceptors, 5
Blanchard, J.C., see Castel, M.N., 111
Blennow, K., see Prior, R., 69
Bogucki, A., see Guiroy, D.C., 31
Bonuccelli, U., Garant, D. and Fariello, R., The acute convulsant effect of MPTP is dependent on intracerebral MPP, 22
Bourgoin, S., see Benoliel, J.J., 204
Brady, R.J., Smith, K.L. and Swann, J.W., Calcium modulation of the *N*-methyl-D-aspartate (NMDA) response and electrographic seizures in immature hippocampus, 92

Cahusac, P.M.B., Rolls, E.T. and Marriott, F.H.C., Potentiation of neuronal responses to natural visual input paired with postsynaptic activation in the hippocampus of the awake monkey, 39
Castel, M.N., Lechardeur, D., Blanchard, J.C. and Laduron, P.M., Reduction of the amount of neurotensin retrogradely transported in dopaminergic neurons of senescent rats, 111
Cecconi, R., see Tramontana, M., 79
Cesselin, F., see Benoliel, J.J., 204
Cherubini, E., see Ito, S., 157
Church, J., Shacklock, J.A. and Baimbridge, K.G., Dextromethorphan and phencyclidine receptor ligands: Differential effects on K- and NMDA-evoked increases in cytosolic free Ca^{2+} concentration, 232
Coenen, T., see Tensen, C.P., 178
Coleman, R.A., see Birrell, G.J., 5
Cools, A.R., see Van den Bos, R., 246
Corio, M., see Portal, J.-J., 105
Cuber, J.-C., see Kitabgi, P., 9

Dani, J.A., see Amador, M., 251
Deecke, L., see Uhl, F., 256
Delalle, I., see Kostović, I., 153
Delebassee, D., see Laurens, C., 1
Delrée, P., see Martin, Didier, 44
Djamgoz, M.B.A., see Fraser, S.P., 242
Duron, Bernard and Khater-Boidin, J., The post-potential excitability modulation of α nerve and muscle fibers in human newborn and adult, 122

Edwards, D.M., see Fry, K.R., 216

Fariello, R., see Bonuccelli, U., 22
Faßbender, C.-P., see Löscher, W., 115
Fenger, M., Processing of pro-opiomelanocortin-derived amidated joining peptideand glycine-extended precursor in monkey pituitary, 190
Ferroni, S., see Schubert, P., 174
Foehring, R.C. and Waters, R.S., Contributions of low-threshold calcium current and anomalous rectifier (I_h) to slow depolarizations underlying burst firing in human neocortical neurons in vitro, 17
Fournier, J., see Laurens, C., 1
Franzen, P., see Uhl, F., 256
Fraser, S.P., Barrett, P., Djamgoz, M.B.A. and Morgan, P.J., Melatonin receptor mRNA expression in *Xenopus* oocytes: inhibition of G-protein-activated response, 242
Frehel, D., see Laurens, C., 1
Fry, K.R., Edwards, D.M., Shaw, K.A. and Watt, C.B., The rabbit retina: a long-term model system for aluminum-induced neurofibrillary degeneration, 216
Frysztak, R.J., see Gonzalez-Lima, F., 74
Fukami, Y., see Onozuka, M., 35
Furuichi, H., see Onozuka, M., 35

Garant, D., see Bonuccelli, U., 22
Gatzinsky, K.P., Berthold, C.-H. and Rydmark, M., Axon–Schwann cell networks are regular components of nodal regions in normal large nerve fibres of cat spinal roots, 264
Gauthier, T., see Laurens, C., 1
Geppetti, P., see Tramontana, M., 79
Gill, B.K., see Nakamura-Craig, M., 49
Gillardon, F., Morano, I. and Zimmermann, M., Ultraviolet irradiation of the skin attenuates calcitonin gene-related peptide mRNA expression in rat dorsal root ganglion cells, 144
Gonzalez-Lima, F. and Frysztak, R.J., Functional mapping of the rat brain during vocalizations: a 2-deoxyglucose study, 74
Gordon, D., see Moskowitz, H., 148
Gottfries, C.G., see Prior, R., 69
Gramsbergen, A. and IJkema-Paassen, J., Increased cell number in remaining cerebellar nuclei after cerebellar hemispherectomy in neonatal rats, 97
Grubb, B.D., see Birrell, G.J., 5
Grünwald, F., Schröck, H. and Kuschinsky, W., The influence of

nicotine on local cerebral blood flow in rats, 108
Guiroy, D.C., Bogucki, A., Papierz, W., Liberski, P.P. and Yanagihara, R., Amyloid β-protein in cerebral amyloid angiopathy, senile plaques, and preamyloidotic lesions in subcortical arteriosclerotic encephalopathy (Binswanger disease), 31

Haass, M., see Hänze, J., 119
Hamon, M., see Benoliel, J.J., 204
Hansen, G.H., Belhage, B. and Schousboe, A., Effect of a GABA agonist on the expression and distribution of $GABA_A$ receptors in the plasma membrane of cultured cerebellar granule cells: an immunocytochemical study, 162
Hänze, J., Kummer, W., Haass, M. and Lang, R.E., Neuropeptide Y mRNA regulation in rat sympathetic ganglia: effect of reserpine, 119
Harrison, N.L., see Lambert, N.A., 101
Hartschuh, W., see Weihe, E., 260
Hatta, K., Yamamato, T., Hori, T., Okuwa, M. and Moroji, T., 3-((±)2-Carboxypiperazin-4-yl)propyl-1-phosphonic acid (CPP) more potently antagonizes the high-affinity Mg^{2+} binding site on the *N*-methyl-D-aspartate/phencyclidine receptor ion channel complex than the L-glutamate recognition site, 229
Hefter, H., see Hummelsheim, H., 52
Hönack, D., see Löscher, W., 115
Hoppe, E., see Wasserkort, R., 183
Hori, T., see Hatta, K., 229
Houpt, T.A., see Mistlberger, R.E., 27
Howes, K.A., see Lerchl, A., 213
Hummelsheim, H. and Hefter, H., A premovement silent period does not occur prior to rapid changes of velocity during human limb movements, 52
Hwa, G.G.C. and Avoli, M., Hyperpolarizing inward rectification in rat neocortical neurons located in the superficial layers, 65
Hyodo, S., see Matsumoto, A., 225

Iggo, A., see Birrell, G.J., 5
Imai, S., see Onozuka, M., 35
Ishihara, A., Taguchi, S., Araki, H. and Nishihira, Y., Retrograde neuronal labeling of motoneurons in the rat by fluorescent tracers, and quantitative analysis of oxidative enzyme activity in labeled neurons, 141
Itano, Y., Noji, S., Koyama, E., Taniguchi, S., Taga, N., Takahashi, T., Ono, K. and Kosaka, F., Bacterial endotoxin-induced expression of metallothionein genes in rat brain, as revealed by in situ hybridization, 13
Ito, S., Provini, L. and Cherubini, E., L-Homocysteic acid mediates synaptic excitation at NMDA receptorsin the hippocampus, 157
Iwata, H., see Koyama, Y., 235

Jarrott, B., see Beart, P.M., 187
Jean, A., see Tell, F., 221
Jukić, S., see Kostović, I., 153
Jung, Daniel, Pons, F., Legér, J.J., Aunis, D. and Rendon, A., Dystrophin in central nervous system: a developmental, regional distribution and subcellular localization study, 87

Kaiya, H., see Nishida, A., 133
Kajander, K.C., see Wakisaka, S., 200
Kanaho, Ya., see Nishida, A., 133
Khater-Boidin, J., see Duron, B., 122
Kim, S.U., see Mizuguchi, M., 166
Kitabgi, P., Masuo, Y., Nicot, A., Berod, A., Cuber, J.-C. and Rostène, W., Marked variations of the relative distributions of neurotensin and neuromedin N in micropunched rat brain areas suggest differential processing of their common precursor, 9
Kosaka, F., see Itano, Y., 13
Kostović, I., Štefulj-Fučić, A., Mrzljak, L., Jukić, S. and Delalle, I., Prenatal and perinatal development of the somatostatin-immunoreactive neurons in the human prefrontal cortex, 153
Koyama, E., see Itano, Y., 13
Koyama, Y., Sugimoto, T., Shigenaga, Y., Baba, A. and Iwata, H., A morphological study on glutamate-induced swelling of cultured astrocytes: involvement of calcium and chloride ion mechanisms, 235
Kummer, W., see Hänze, J., 119
Kuschinsky, W., see Grünwald, F., 108

Laduron, P.M., see Castel, M.N., 111
Lambert, N.A., Harrison, N.L. and Teyler, T.J., Evidence for μ opiate receptors on inhibitory terminals in area CA1 of rat hippocampus, 101
Lang, R.E., see Hänze, J., 119
Lang, W., see Uhl, F., 256
Laurens, C., Fournier, J., Delebassee, D., Frehel, D., Gauthier, T. and Soubrie, P., Enhancement of neurite outgrowth from central nervous system neurons in primary culture by thrombin inhibitors, 1
Lechardeur, D., see Castel, M.N., 111
Legér, J.J., see Jung, D., 87
Legrand, J.C., see Benoliel, J.J., 204
Lemos, J.R., see Wang, X., 125
Leprince, P., see Martin, D., 44
Lerchl, A., Reiter, R.J., Howes, K.A., Nonaka, K.O. and Stokkan, K.-A., Evidence that extremely low frequency Ca^{2+}-cyclotron resonance depresses pineal melatonin synthesis in vitro, 213
Liberski, P.P., see Guiroy, D.C., 31
Lindinger, G., see Uhl, F., 256
Löscher, W., Hönack, D. and Faßbender, C.-P., Regional alterations in brain amino acids after administration of the *N*-methyl-D-aspartate receptor antagonists MK-801 and CGP 39551 in rats, 115

Mager, R., see Schubert, P., 174
Maggi, C.A., see Tramontana, M., 79
Marcus, J.T. and Toet, A., Spatiotemporal representation of moving luminance edges in human vision, 239
Marriott, F.H.C., see Cahusac, P.M.B., 39
Martin, D., Schoenen, J., Delrée, P., Leprince, P., Rogister, B. and Moonen, G., Grafts of syngenic cultured, adult dorsal root ganglion-derived Schwann cells to the injured spinal cord of adult rats: preliminary morphological studies, 44
Masters, C.L., see Prior, R., 69
Masuo, Y., see Kitabgi, P., 9
Matsumoto, A., Arai, Y., Uran, A., and Hyodo, S., Cellular localization of gap junction mRNA in the neonatal rat brain, 225
Mauborgne, A., see Benoliel, J.J., 204
McQueen, D.S., see Birrell, G.J., 5
Medvedev, S.V., see Bechtereva, N.P., 61
Mercer, L.D., see Beart, P.M., 187
Mistlberger, R.E., Houpt, T.A. and Moore-Ede, M.C., The benzodiazepine triazolam phase-shifts circadian activity rhythms in a diurnal primate, the squirrel monkey (*Saimiri sciureus*), 27
Mitchell, J.J. and Anderson, K.J., Quantitative autoradiographic analysis of excitatory amino acid receptors in the cat spinal cord, 269
Mizuguchi, M. and Kim, S.U., α_1-Antichymotrypsin supports short-term survival of cerebral neurons in culture, 166
Møller, A.R., see Schmid, U.D., 273
Mönning, U., see Prior, R., 69

Moonen, G., see Martin, D., 44
Moore-Ede, M.C., see Mistlberger, R.E., 27
Morano, I., see Gillardon, F., 144
Morgan, P.J., see Fraser, S.P., 242
Moroji, T., see Hatta, K., 229
Morris, M.E., see Rosen, A.S., 169
Moskowitz, H., Zlotkin, E. and Gordon, D., Solubilization and characterization of the insect neuronal sodium channel, 148
Mrzljak, L., see Kostović, I., 153

Nakai, Ya., see Wang, QP., 129
Nakamura-Craig, M. and Gill, B.K., Effect of neurokinin A, substance P and calcitonin gene related peptide in peripheral hyperalgesia in the rat paw, 49
Nicot, A., see Kitabgi, P., 9
Nishida, A., Kaiya, H., Kanaho, Ya. and Nozawa, Y., Distinct cellular expression of pertussis toxin-sensitive GTP-binding proteins in rat cerebellum, 133
Nishihira, Y., see Ishihara, A., 141
Nohr, D., see Weihe, E., 260
Noji, S., see Itano, Y., 13
Nonaka, K.O., see Lerchl, A., 213
Nozawa, Y., see Nishida, A., 133

Okuwa, M., see Hatta, K., 229
Ono, K., see Itano, Y., 13
Onozuka, M., Furuichi, H., Imai, S. and Fukami, Y., Evidence that Ca^{2+}/calmodulin-dependent protein phosphorylation is involved in the opening process of potassium channels in identified snail neurons, 35
Osipenko, O., see Belan, P.V., 137

Papierz, W., see Guiroy, D.C., 31
Phillis, J.W., see Simpson, R.E., 83
Pons, F., see Jung, D., 87
Portal, J.-J., Corio, M. and Viala, D., Localization of the lumbar pools of motoneurones which provide hindlimb muscles in the rabbit, 105
Prior, R., Mönning, U., Schreiter-Gasser, U., Weidemann, A., Blennow, K., Gottfries, C.G., Masters, C.L. and Beyreuther, K., Quantitative changes in the amyloid βA4 precursor protein in Alzheimer cerebrospinal fluid, 69
Provini, L., see Ito, S., 157

Randić, M., see Rusin, K.I., 208
Reddington, M., see Wasserkort, R., 183
Reiter, R.J., see Lerchl, A., 213
Rendon, A., see Jung, D., 87
Rogister, B., see Martin, D., 44
Rojas, L. and Zuazaga, C., Spontaneous openings of large and small conductance acetylcholine receptor channels in *Xenopus* myocytes, 195
Rolls, E.T., see Cahusac, P.M.B., 39
Rosen, A.S. and Morris, M.E., Depolarizing effects of anoxia on pyramidal cells of rat neocortex, 169
Rostène, W., see Kitabgi, P., 9
Rusin, K.I. and Randić, M., Modulation of NMDA-induced currents by μ-opioid receptor agonist DAGO in acutely isolated rat spinal dorsal horn neurons, 208
Rydmark, M., see Gatzinsky, K.P., 264

Santicioli, P., see Tramontana, M., 79
Schmid, J., see Schmid, U.D., 273
Schmid, U.D., Møller, A.R. and Schmid, J., Transcranial magnetic stimulation excites the labyrinthine segment of the facial nerve: an intraoperative electrophysiological study in man, 273
Schoenen, J., see Martin, D., 44
Schousboe, A., see Hansen, G.H., 162
Schreiter-Gasser, U., see Prior, R., 69
Schröck, H., see Grünwald, F., 108
Schubert, P., Ferroni, S. and Mager, R., Pharmacological blockade of Cl^- pumps or Cl^- channels reduces the adenosine-mediated depression of stimulus train-evoked Ca^{2+} fluxes in rat hippocampal slices, 174
Schubert, P., see Wasserkort, R., 183
Shacklock, J.A., see Church, J., 232
Shaw, K.A., see Fry, K.R., 216
Shigenaga, Y., see Koyama, Yu., 235
Shioda, S., see Wang, QP., 129
Simpson, R.E., Walter, G.A. and Phillis, J.W., The effects of hypothermia on amino acid neurotransmitter release from the cerebral cortex, 83
Smith, K.L., see Brady, R.J., 92
Soubrie, P., see Laurens, C., 1
Štefulj-Fučić, A., see Kostović, I., 153
Stemple, D.L. and Anderson, D.J., A Schwann cell antigen recognized by monoclonal antibody 217c is the rat low-affinity nerve growth factor receptor, 57
Stokkan, K.-A., see Lerchl, A., 213
Sugimoto, T., see Koyama, Yu., 235
Swann, J.W., see Brady, R.J., 92

Taga, N., see Itano, Y., 13
Taguchi, S., see Ishihara, A., 141
Takahashi, T., see Itano, Y., 13
Taniguchi, S., see Itano, Y., 13
Tell, F. and Jean, A., Bursting discharges evoked in vitro,by solitary tract stimulation or application of *N*-methyl-D-aspartate,in neurons of the rat nucleus tractus solitarii, 221
Tensen, C.P., Coenen, T. and Van Herp, F., Detection of mRNA encoding Crustacean Hyperglycemic Hormone (CHH) in the eyestalk of the crayfish *Orconectes limosus* using non-radioactive in situ hybridization, 178
Teyler, T.J., see Lambert, N.A., 101
Toet, A., see Marcus, J.T., 239
Tramontana, M., Cecconi, R., Bianco, E.D., Santicioli, P., Maggi, C.A., Alessandri, Massimo and Geppetti, P., Hypertonic media produce Ca^{2+}-dependent release of calcitonin gene-related peptide from capsaicin-sensitive nerve fibres in the rat urinary bladder, 79
Treistman, S.N., see Wang, X., 125

Uhl, F., Franzen, P., Lindinger, G., Lang, W. and Deecke, L., On the functionality of the visually deprived occipital cortex in early blind persons, 256
Urano, A., see Matsumoto, A., 225

Van den Bos, R. and Cools, A.R., Motor activity and the $GABA_A$-receptor in the ventral pallidum/substantia innominata complex, 246
Van Herp, F., see Tensen, C.P., 178
Viala, D., see Portal, J.-J., 105

Wakisaka, S., Kajander, K.C. and Bennett, G.J., Increased neuropeptide Y (NPY)-like immunoreactivity in rat sensory neurons following peripheral axotomy, 200

Walter, G.A., see Simpson, R.E., 83
Wang, QP., Nakai, Y. and Shioda, S., Enkephalinergic synaptic axon terminals on serotoninergic neurons in the dorsal raphe nucleus of the rat: an electron microscopic study by the double immunostaining method, 129
Wang, X., Treistman, S.N. and Lemos, J.R., Direct identification of individual vasopressin-containing nerve terminals of the rat neurohypophysis after 'whole-cell' patch-clamp recordings, 125
Wasserkort, R., Hoppe, E., Reddington, M. and Schubert, P., Modulation of A_1 adenosine receptor function in rat brain by the polyamine, spermine, 183
Waters, R.S., see Foehring, R.C., 17
Watt, C.B., see Fry, K.R., 216
Weidemann, A., see Prior, R., 69
Weihe, E., Hartschuh, W. and Nohr, D., Light microscopic immunoenzyme and electron microscopic immunogold cytochemistry reveal tachykinin immunoreactivity in Merkel cells of pig skin, 260
Ykema-Paassen, J., see Gramsbergen, A., 97
Yamamato, T., see Hatta, K., 229
Yanagihara, R., see Guiroy, D.C., 31
Zimmermann, M., see Gillardon, F., 144
Zlotkin, E., see Moskowitz, H., 148
Zuazaga, C., see Rojas, L., 195

ANNOUNCEMENT

The *3rd International Congress of the International Society of Neuroimmunology* will be held in Jerusalem, Israel, from October 27 to November 1, 1991.

For further information: Secretariat ISNI
P.O. Box 50006
61500 Tel Aviv
Israel
Tel. 972 3 654571
Fax 972 3 655674
Tlx. 341171 KENS IL

CONTENTS *(continued from back cover)*

Dextromethorphan and phencyclidine receptor ligands: Differential effects on K^+- and NMDA-evoked increases in cytosolic free Ca^{2+} concentration
J. Church, J.A. Shacklock and K.G. Baimbridge (Canada) 232
A morphological study on glutamate-induced swelling of cultured astrocytes: involvement of calcium and chloride ion mechanisms
Y. Koyama, T. Sugimoto, Y. Shigenaga, A. Baba and H. Iwata (Japan) 235
Spatiotemporal representation of moving luminance edges in human vision
J.T. Marcus and A. Toet (The Netherlands) 239
Melatonin receptor mRNA expression in *Xenopus* oocytes: inhibition of G-protein-activated response
S.P. Fraser, P. Barrett, M.B.A. Djamgoz and P.J. Morgan (U.K.) 242
Motor activity and the $GABA_A$-receptor in the ventral pallidum/substantia innominata complex
R. van den Bos and A.R. Cools (The Netherlands) 246
Protein kinase inhibitor, H-7, directly affects *N*-methyl-D-aspartate receptor channels
M. Amador and J.A. Dani (U.S.A.) 251
On the functionality of the visually deprived occipital cortex in early blind persons
F. Uhl, P. Franzen, G. Lindinger, W. Lang and L. Deecke (Austria) 256
Light microscopic immunoenzyme and electron microscopic immunogold cytochemistry reveal tachykinin immunoreactivity in Merkel cells of pig skin
E. Weihe, W. Hartschuh and D. Nohr (F.R.G.) 260
Axon–Schwann cell networks are regular components of nodal regions in normal large nerve fibres of cat spinal roots
K.P. Gatzinsky, C.-H. Berthold and M. Rydmark (Sweden) 264
Quantitative autoradiographic analysis of excitatory amino acid receptors in the cat spinal cord
J.J. Mitchell and K.J. Anderson (U.S.A.) 269
Transcranial magnetic stimulation excites the labyrinthine segment of the facial nerve: an intraoperative electrophysiological study in man
U.D. Schmid, A.R. Møller and J. Schmid (U.S.A.) 273